



DOI: <http://dx.doi.org/10.1590/1807-1929/agriambi.v21n12p817-821>

## Spikelet sterility in rice genotypes affected by temperature at microsporogenesis

Natalia M. de Souza<sup>1</sup>, Rubens Marschalek<sup>2</sup>, Luis Sangoi<sup>1</sup> & Francieli S. Weber<sup>1</sup>

<sup>1</sup> Universidade do Estado de Santa Catarina/Centro de Ciências Agroveterinárias. Lages, SC. E-mail: [naty\\_natynatalia@hotmail.com](mailto:naty_natynatalia@hotmail.com) (Corresponding author); [luis.sangoi@udesc.br](mailto:luis.sangoi@udesc.br); [francieliweber@yahoo.com.br](mailto:francieliweber@yahoo.com.br)

<sup>2</sup> Epagri/Estação Experimental de Itajaí. Itajaí, SC. E-mail: [rubensm@epagri.sc.gov.br](mailto:rubensm@epagri.sc.gov.br)

### Key words:

*Oryza sativa*  
booting  
cold  
controlled environment

### ABSTRACT

This study evaluated the effect of temperatures during the phase of microsporogenesis on spikelet sterility of paddy rice and identified genotypes tolerant to low temperatures at this growth stage. The inbreds SC681, SC491, and SC676 and the cultivars Epagri 109 and SCS116 Satoru were assessed. The genotypes were submitted for three days in a growth chamber to five temperatures at microsporogenesis: 9, 12, 15, 18, and 21 °C. For each tested temperature, a control was kept in the greenhouse under environmental conditions. After harvest, full and empty spikelets were counted and weighed and the percentage of spikelet sterility was determined. Data were evaluated by variance analysis using the F test. Averages were compared by Tukey's test and regression analysis. The highest spikelet sterilities were observed when the genotypes were exposed to the temperatures of 9 and 12 °C. Genotype spikelet sterility was similar to that of the control at 21 °C. The inbred SC 676 presented higher tolerance to lower temperatures is therefore potentially suited to generate a cultivar with adequate agronomic performance in rice growing areas prone to cold conditions at microsporogenesis.

### Palavras-chave:

*Oryza sativa*  
emborrachamento  
frio  
ambiente controlado

## Esterilidade de espiguetas em genótipos de arroz irrigado afetada pela temperatura na microsporogênese

### RESUMO

Este trabalho teve como objetivo avaliar o efeito da temperatura na fase de microsporogênese sobre a esterilidade de espiguetas de arroz irrigado identificando, nesta fase, genótipos tolerantes a baixas temperaturas. Foram testadas as linhagens SC681, SC491 e SC676 e as cultivares Epagri 109 e SCS116 Satoru. Os genótipos foram submetidos por três dias em câmara de crescimento a cinco temperaturas na microsporogênese: 9, 12, 15, 18 e 21 °C. Para cada temperatura um tratamento testemunha foi mantido a temperatura ambiente na casa de vegetação. Após colheita realizaram-se a contagem e a pesagem das espiguetas cheias e vazias, determinando-se a percentagem de esterilidade. Os dados foram submetidos à análise de variância pelo teste F. Quando significativas, as médias foram comparadas pelo teste de Tukey e análise de regressão. A maior esterilidade foi observada nos genótipos submetidos às temperaturas de 9 e 12 °C. A esterilidade foi similar entre testemunhas e a 21 °C. A linhagem SC676 apresentou maior tolerância à ocorrência de baixas temperaturas mostrando-se promissora para gerar uma cultivar com desempenho agrônômico adequado, em regiões orizícolas propensas à ocorrência de frio na microsporogênese.



## INTRODUCTION

Paddy rice is grown in several regions, climatic conditions, and production systems around the world (Artacho et al., 2011). However, low temperatures during critical growth stages affect its productivity (Shimono et al., 2007; SOSBAI, 2014). The extension of damage caused by cold incidence depends on several factors, such as the duration and intensity of thermic stress, crop management, cultivar, and plant development stage (Cruz & Milach, 2000).

The growth stage R2 (microsporogenesis) of the scale proposed by Counce et al. (2000) is the phase most sensitive to low temperatures (Rozzetto et al., 2013). At this stage, temperatures below 17 °C negatively impact pollen grain formation (SOSBAI, 2014), resulting in increased spikelet sterility in this period (Baruah et al., 2009).

In the South of Brazil, where the two larger paddy rice producer states are located, Rio Grande do Sul and Santa Catarina, low temperatures are one of the major factors reducing crop grain yields (Cruz et al., 2006). Rice is a spring/summer cereal that can be injured by cold throughout the entire growth cycle, from germination to grain maturity, resulting in reduced productivity (Shimono et al., 2007). Cold tolerance during the reproductive period is important to assure high yields in environments where low temperatures are common (Cruz et al., 2010).

In this context, this work evaluated the effect of temperature at microsporogenesis on spikelet sterility of paddy rice, with the aim to identify genotypes tolerant to low temperatures at this growth stage.

## MATERIAL AND METHODS

The study was carried out at the experimental station of Epagri, located in Itajaí, SC, during the growing season of 2013/2014. The experiment was set up in recipients filled with Dystric Gleysol. Each recipients corresponded to one experimental unit. The recipients were placed in a greenhouse and a growth chamber. The soil chemical attributes were assessed according to the methodology described by Tedesco et al. (1995), obtaining the following values: pH-H<sub>2</sub>O = 5.1; P = 11.9 mg kg<sup>-1</sup>; K = 64 mg kg<sup>-1</sup>; MO = 1.7%; Al = 2.9 cmol<sub>c</sub> dm<sup>-3</sup>; Ca = 0.3 cmol<sub>c</sub> dm<sup>-3</sup>; Mg = 0.4 cmol<sub>c</sub> dm<sup>-3</sup>, and clay content = 230 g dm<sup>-3</sup>. The experimental design was completely randomized, with treatments disposed in a multifactorial arrangement (5 x 5 x 2) with three replicates. The first factor corresponded to five late ripening genotypes: three inbred lines (SC681, SC491, and SC676) and two commercial cultivars (Epagri 109 and SCS116 Satoru). The inbred lines were selected based on their promising behavior when submitted to low temperatures in experiments carried out by Marschalek et al. (2013) in the High Valley of Itajaí, an important rice production region in the state of Santa Catarina. The selected cultivars are extensively cultivated in Santa Catarina and sensitive to cold weather, previously shown by SCS116 Satoru. During microsporogenesis, each genotype was placed in a growth chamber and submitted to five temperatures: 9, 12, 15, 18, and 21 °C, corresponding to the second factor. For each evaluated temperature and genotype,

a control was kept in the greenhouse under environmental temperatures, corresponding to the third factor. This procedure was needed because the growth chamber only had the capacity to accommodate one thermic treatment per time. Therefore, the five temperature regimes had to be split and the experiment was composed of 150 experimental units (5 x 5 x 2 x 3).

The experimental units had the following dimensions: 22 cm of diameter, 20 cm of height, and the capacity to store 7 kg of soil. In each recipients, 10 to 15 seeds were sown. After seedling emergence, when plants were at the V2 growth stage, the first thinning was performed, leaving four plants per bucket. At the V6 growth stage, a second thinning was carried out, assuring a final population of two plants in each experimental unit.

All treatments received the same fertilization rates and products, according to the recommendations of SOSBAI (2014) and the results of the soil analysis. The following fertilizer quantities were applied per recipients: 2 g of triple superphosphate, corresponding to P<sub>2</sub>O<sub>5</sub> fertilization, and 20 mL of a N + K<sub>2</sub>O solution on each top-dressing fertilization. This solution was obtained through the addition and agitation of urea (200 g) and potassium chloride (160 g), diluted in 2 L of water. Application rates were equivalent to fertilization with 200 mg kg<sup>-1</sup> N, 120 mg kg<sup>-1</sup> P<sub>2</sub>O<sub>5</sub>, and 70 mg kg<sup>-1</sup> K<sub>2</sub>O.

The recipients remained in the greenhouse from sowing (stage S0) to microsporogenesis (stage R2). Six tillers were tagged when they reached R2; this growth stage was identified following the plant development on each experimental unit daily. The tiller stems were periodically opened until the booting stage (R2) was observed.

According to Zaffari et al. (2014), rice microsporogenesis can be identified based on the distance between the flag leaf ligule and the penultimate leaf ligule. This distance must be comprised between 3 cm (flag leaf ligule 3 cm below penultimate leaf ligule) and 10 cm (flag leaf ligule 10 cm above penultimate leaf ligule). Since plant tillers do not reach the booting stage at the same time, tillers were marked when the distance between the last two leaf ligules ranged from 1 to 2 cm.

After tillers were tagged, the recipients were transferred to the growth chamber so that each thermic regime could be applied. Plants were submitted for three consecutive days to one of the five temperatures previously determined by the trial protocol while the control plants remained in the greenhouse. During this period, light and air relative humidity (RH) in the growth chamber were monitored and kept at 12 h of light, 12 h of dark, and 65% of RH. After three days, the recipients were returned to the greenhouse and kept until harvest.

At the end of the crop cycle, tagged panicles were individually harvested. The remaining panicles from each recipients were harvested in bulk. Each genotype was manually shelled. Empty and full spikelets were separated with a fan and counted. Full and empty spikelet weight and sterility percentage of individual and bulk panicles were determined.

The data were submitted to a variance analysis using the F test at the significance level of 0.05 probability. When the F values were significant, the means were compared by Tukey Test. In cases where the temperature effect was significant, a polynomial regression analysis was also performed. Both mean comparisons were carried out at the significance level of 0.05

probability. The data were analyzed with the Assistat program (Silva & Azevedo, 2016).

## RESULTS AND DISCUSSION

Variance analysis for the variable “bulk spikelet sterility” detected as significant the triple interaction among genotype x temperature x thermic stress/control, indicating that each genotype responded differently to the temperature variation and the imposition of the thermic stress (Table 1).

All genotypes presented a similar percentage of bulk sterility in the control (without application of thermic stress) and when submitted to 21 °C at microsporogenesis (Table 2). This indicates that the temperature of 21 °C did not interfere negatively with grain pollen formation. Therefore, the genotypes presented the same behavior compared to the treatment without thermic stress. The rate of genotype spikelet sterility ranged from 25.8 to 36.6% at the temperature of 21 °C and in the control. These results differed from those gathered by Rozetto et al. (2013), who observed lower percentages of spikelet sterility in the control, varying between 7.8 and 19.7%, in an experiment carried out under the same conditions.

The temperature of 9 °C had the greatest impact on microsporogenesis of the evaluated genotypes, causing sterility levels of 100% on Epagri 109, SC491, and SC681, 43.3% on SC676, and 33.5% on SCS116 Satoru (Table 2). The lowest percentage of spikelet sterility expressed by cultivar SCS116 Satoru at 9 °C did not agree with the findings of previous studies because this cultivar was included in the trial due to its cold sensitivity, reported in experiments carried out by Marschalek et al. (2013) in the High Valley of Itajai. The inbred SC676 presented the lowest numeric value of spikelet sterility at the temperature of 18 °C. The genotype's percentage of bulk spikelet sterility did not differ significantly between the temperatures 15 and 21 °C.

Regression analysis of the interaction between temperature and bulk spikelet sterility showed a quadratic behavior of the evaluated genotypes, with the exception of cultivar SCS116 Satoru, for which the regression was not significant (Figure

Table 1. Variance analysis for the variables bulk and individual panicle spikelet sterility of paddy rice

Source of variation	D.F.	S.Q.	M.S.	F
Bulks				
Genotype	4	1,524.86	381.21	4.55*
Temperature	4	7,458.03	1,864.51	22.24*
Thermic Stress/Control	1	1,1867.30	11,867.30	141.58*
Genotype x Temperature	16	5,937.35	371.08	4.43*
Genotype x Thermic stress/Control	4	4,031.55	1,007.89	12.02*
Temperature x Thermic stress/Control	4	17,391.70	4,347.92	51.87*
Genotype x Temp x Thermic stress/Control	16	6,666.94	416.68	4.97*
Treatments	49	54,877.70	1,119.95	13.36*
Residue	100	8,381.71	83.82	
Total	149	63,259.40		
Individual Panicles				
Genotype	4	1,304.70	326.17	2.14 <sup>NS</sup>
Temperature	4	16,770.00	4,192.51	27.46*
Thermic Stress/Control	1	25,545.40	25,545.40	167.34*
Genotype x Temperature	16	1,719.24	107.45	0.70 <sup>NS</sup>
Genotype x Thermic stress/Control	4	3,772.59	943.15	6.18*
Temperature x Thermic stress/Control	4	33,887.90	8,471.98	55.50*
Genotype x Temp x Thermic stress/Control	16	2,403.09	150.19	0.98 <sup>NS</sup>
Treatment	49	8,5403.00	1,742.92	11.42*
Residue	100	15,265.20	152.65	
Total	149	100,668.00		

\*Significant at the probability level of 0.05; NS – Not significant at the probability level of 0.05

Table 2. Percentage of bulk spikelet sterility in five paddy rice genotypes submitted to five temperatures at microsporogenesis, in comparison to the control

Thermic regime	Epagri 109	SC491	SC676	SC681	SCS 116 Satoru
Bulk spikelet sterility (%)					
9 °C	100.0 aA*	100.0 aA	43.3 bA	100.0 aA	33.5 bABC
Control	13.2 aC	18.3 aCD	24.6 aAB	18.7 aCD	14.3 aC
12 °C	22.1 bBC	47.9 aB	38.5 abAB	35.8 abBCD	39.5 abAB
Control	18.0 aBC	14.3 aD	17.2 aB	16.5 aD	13.8 aC
15 °C	31.0 aBC	39.5 aBC	27.0 aAB	40.8 aBC	47.5 aA
Control	30.6 aBC	19.9 aCD	31.2 aAB	22.7 aBCD	20.6 aBC
18 °C	29.4 abBC	41.3 aBC	19.6 aAB	45.1 aB	33.7 abABC
Control	39.6 aB	36.0 aBCD	41.3 aAB	31.1 aBCD	35.3 aABC
21 °C	28.2 aBC	34.2 aBCD	25.8 aAB	29.6 aBCD	31.9 aABC
Control	27.4 aBC	29.0 aBCD	36.6 aAB	22.3 aBCD	28.0 aA
CV% = 27.15					

\*Means followed by the same lowercase letter in the row and uppercase letter in the column are not significantly different by Tukey Test at the significance level of 0.05 probability

1). According to the equations adjusted to the data, the smallest values for this variable were located between 17 and 18 °C, corroborating the observations made by Walter et al. (2010), Nanculao et al. (2013) and Oort et al. (2014), and that temperatures below these values increase rice spikelet sterility. The inbred SC676 presented the smallest rate of increment in bulk spikelet sterility when temperatures dropped from 21 to 9 °C.

The sterility of panicles harvested separately was affected by the double interactions between genotype x thermic stress/control and temperature x thermic stress/control, indicating that in the presence of thermic stress, genotypes and temperature responded differently from the controls.

Considering the tagged panicles, the five genotypes had similar percentages of spikelet sterility on the controls that were not subjected to thermic stress (Table 3). Conversely, in the




Figure 1. Effect of temperature on the percentage of bulk spikelet sterility of five paddy rice genotypes

Table 3. Overall percentage of tagged panicle spikelet sterility of five paddy rice genotypes in comparison to the control averaged across all five temperatures

Thermic regime	Epagri 109	SC491	SC676	SC681	SCS116 Satoru
Tagged panicle spikelet sterility (%)					
Temperatures <sup>1/</sup>	43.0 bcA*	59.5 aA	39.7 aA	54.9 abA	55.3 abA
Control	24.1 aB	23.7 aB	29.8 aB	21.0 Ab	23.2 aB
CV% = 33.02					

\* Means followed by the same lowercase letter in the row and uppercase letter in the column are not significantly different by Tukey Test at the significance level of 0.05 probability

<sup>1/</sup> Average values of five temperatures: 9, 12, 15, 18, and 21 °C

presence of thermic stress, the highest sterilities were registered in the inbreds SC491 and SC681 and in the cultivar SCS116 Satoru. The inbred SC676 presented the lowest percentage of spikelet sterility at all five temperatures. All genotypes had larger sterility than the control when exposed to thermic stress. Nonetheless, the smallest numeric difference in the percentage of marked panicle spikelet sterility between stressed plants and control was registered for the inbred SC676.

Across all five genotypes, there was a quadratic decrease in the percentage of tagged panicle spikelet sterility when temperatures increased from 9 to 20 °C, demonstrating that the smallest sterility percentages were obtained at the higher temperatures assessed in the experiment (Figure 2). This behavior reinforced the observations made by Khan et al. (1986), Kaw (1991), Streck et al. (2006), Baruah et al. (2009), Wang et al. (2009), Peyman & Hashem (2010), and Shinada et al. (2013), emphasizing that extremely low temperatures increase rice spikelet sterility and temperatures above 20 °C do not compromise grain pollen formation at microsporogenesis. Considering cold-tolerant genotypes, the critical temperatures to trigger spikelet sterility ranged from 15 to 17 °C, whereas for cold-sensitive genotypes, it was between 17 and 19 °C (SOSBAI, 2014). This information was confirmed in the present study where the greatest sterility rates were detected when plants were submitted to 9, 12, and 15 °C during microsporogenesis (Table 2 and Figures 1 and 2). On the other hand, when temperatures varied from 18 to 21 °C, plant behavior was similar to that in the control.

Higher sterility rates caused by the exposition to low temperatures have also been reported by Soltani et al. (2001), Farrell et al. (2006), Martins et al. (2007), Jagadish et al. (2008), Weerakoon et al. (2008), Walter et al. (2010), Wang et al. (2013), and Oort et al. (2014). These studies showed that the stress duration is an important factor as a temperature of 12 °C may not induce sterility if it occurs over a period shorter than 48 h, but can cause 100% of sterility when plants are exposed for more than six days to this stress, depending on the genotype sensitivity.




Figure 2. Effect of temperature on the percentage of tagged panicle spikelet sterility, averaged across the five paddy rice genotypes

One of the main objectives of this work was to identify genotypes with high tolerance to low temperatures, allowing cultivation in regions prone to cold periods during rice microsporogenesis. The spikelet sterility data collected in the experiment demonstrated that the inbred SC676 was more suitable than the other genotypes in terms of tolerating low temperatures at rice booting. It was the only inbred for which bulks did not present 100% sterility when submitted to 9 °C (Table 2). Furthermore, this genotype presented the lowest increment in percentage of spikelet sterility with temperature reduction (Figure 1). Moreover, it also had the lowest rate of tagged panicle sterility under thermic stress, averaged across the five temperatures (Table 3).

Such characteristics indicate that the inbred SC676 is promising to generate a future rice cultivar with favorable agronomic performance in regions with high risks of low temperatures at microsporogenesis. This behavior corroborates the observations made by Marschalek et al. (2013) in field evaluations performed with this inbred at the High Valley of Itajai.

## CONCLUSIONS

1. Temperatures of 9 and 12 °C at microsporogenesis promote high percentage of paddy rice spikelet sterility, regardless of the genotype.
2. Temperatures of 18 and 21 °C do not cause significant damage to paddy rice spikelet sterility.
3. The inbred SC676 presents greater tolerance to low temperatures than the other genotypes, therefore being suitable for the generation of a cultivar with adequate agronomic performance in rice production regions prone to cold stress in the South of Brazil.

## ACKNOWLEDGMENTS

We thank the staff of the Epagri, Itajai Experimental Station, for providing the structure to carry out the experiments. We also thank the UDESC for granting a study scholarship to the first author and CNPq for providing a research scholarship for the second author.

## LITERATURE CITED

- Artacho, P.; Meza, F.; Alcalde, J. A. Evaluation of the ORYZA2000 rice growth model under nitrogen-limited conditions in an irrigated Mediterranean environment. *Chilean Journal of Agricultural Research*, v.71, p.23-33, 2011. <https://doi.org/10.4067/S0718-58392011000100003>
- Baruah, A. R.; Ishigo-Oka, N.; Adachi, M.; Oguma, Y.; Tokizono, Y.; Onishi, K.; Sano, Y. Cold tolerance at the early growth stage in wild and cultivated rice. *Euphytica*, v.165, p.459-470, 2009. <https://doi.org/10.1007/s10681-008-9753-y>
- Counce, P. A.; Keisling, T. C.; Mitchell, A. J. A uniform, objective and adaptive system for expressing rice development. *Crop Science*, v.40, p.436-443, 2000. <https://doi.org/10.2135/cropsci2000.402436x>
- Cruz, R. P. da; Duarte, I. T. de L.; Cabreira, C. Inheritance of rice cold tolerance at the seedling stage. *Scientia Agricola*, v.67, p.669-674, 2010. <https://doi.org/10.1590/S0103-90162010000600008>
- Cruz, R. P. da; Milach, S. C. K. Genetic breeding for cold tolerance in paddy rice. *Ciência Rural*, v.30, p.909-917, 2000.

- Cruz, R. P. da; Milach, S. C. K.; Federizzi, L. C. Rice cold tolerance at the reproductive stage in a controlled environment. *Scientia Agricola*, v.63, p.255-261, 2006. <https://doi.org/10.1590/S0103-90162006000300007>
- Farrell, T. C.; Fox, K. M.; Williams, R. L.; Fukai, S. Genotypic variation for cold tolerance during reproductive development in rice: Screening with cold air and cold water. *Field Crops Research*, v.98, p.178-194, 2006.
- Jagdish, S. V. K.; Craufurd, P. Q.; Wheeler, T. R. Phenotyping parents of mapping populations of rice for heat tolerance during anthesis. *Crop Science*, v.48, p.1140-1146, 2008. <https://doi.org/10.2135/cropsci2007.10.0559>
- Kaw, R. N. Genetic parameters of cold tolerance in rice. *The Indian Journal of Genetics and Plant Breeding*, v.51, p.59-65, 1991.
- Khan, D. R.; Mackill, D. J.; Vergara, B. S. Selection for tolerance to low temperature-induced spikelet sterility at anthesis in rice. *Crop Science*, v.26, p.694-698, 1986. <https://doi.org/10.2135/cropsci1986.0011183X002600040011x>
- Marschalek, R.; Schiocchet, M. A.; Knoblauch, R.; Stuker, H.; Eberhardt, D. S.; Rozzetto, D. S.; Porto, G.; Santos, S. B. dos. Selection of paddy rice genotypes adapted to high land regions prone to low temperatures. In: Congresso Brasileiro de Arroz Irrigado, 8, 2013, Santa Maria. Anais... Santa Maria: Pallotti, 2013. p.181-184.
- Martins, A. F.; Vieira, E. A.; Kopp, M. M.; Luz, V. K. da; Carvalho, M. F. de; Castelo Branco, J. S.; Cruz, R. P. da; Carvalho, F. I. F. de; Oliveira, A. C. de. Characterization of rice families to cold tolerance in the vegetative and reproductive phases. *Bragantia*, v.66, p.227-233, 2007. <https://doi.org/10.1590/S0006-87052007000200006>
- Nanculao, G. D.; Cárcamo, M. P.; Santos, O. A. de los; Velásquez, V. B. Cold tolerance evaluation in Chilean rice genotypes at the germination stage. *Chilean Journal of Agricultural Research*, v.73, p.3-8, 2013. <https://doi.org/10.4067/S0718-58392013000100001>
- Oort, P. A. J. van; Saito, K.; Zwart, S. J.; Shrestha, S. A simple model for simulating heat induced sterility in rice as a function of flowering time and transpirational cooling. *Fields Crops Research*, v.156, p.303-312, 2014. <https://doi.org/10.1016/j.fcr.2013.11.007>
- Peyman, S.; Hashem, A. Evaluation of eighteen rice genotypes in cold tolerance at germination stage. *World Applied Sciences Journal*, v.11, p.1476-1480, 2010.
- Rozzetto, D. S.; Marschalek, R.; Stuker, H.; Eberhardt, D. S.; Raimondi, J. V.; Santos, S. B. dos; Porto, G.; Pazini, B. S.; Souza, N. M. de. Cold tolerance of paddy rice genotypes exposed to low temperatures in growth chamber during the reproductive stage. In: Congresso Brasileiro de Arroz Irrigado, 8, 2013, Anais... Santa Maria: Pallotti, 2013. p.185-188.
- Shimono, H.; Okada, M.; Kanda, E.; Arakawa, I. Low temperature-induced sterility in rice: Evidence for the effects of temperature before panicle initiation. *Field Crops Research*, v.101, p.221-231, 2007. <https://doi.org/10.1016/j.fcr.2006.11.010>
- Shinada, H.; Iwata, N.; Sato, T.; Fujino, K. Genetical and morphological characterization of cold tolerance at fertilization stage in rice. *Breeding Science*, v.63, p.197-204, 2013. <https://doi.org/10.1270/jsbbs.63.197>
- Silva, F. de A. S. e; Azevedo, C. A. V. de. The Assisat Software Version 7.7 and its use in the analysis of experimental data. *African Journal of Agricultural Research*, v.11, p.3733-3740, 2016. <https://doi.org/10.5897/AJAR2016.11522>
- Soltani, A.; Zeinali, E.; Galeshi, S.; Niari, N. Simulating GFDL predicted climate change impacts on rice cropping in Iran. *Journal of Agricultural Science and Technology*, v.3, p.81-90, 2001.
- SOSBAI - Sociedade Sul-Brasileira de Arroz Irrigado. Paddy rice: Research technical recommendations for the South of Brazil. Bento Gonçalves: SOSBAI, 2014. 192p.
- Streck, N. A.; Bosco, L. C.; Michelon, S.; Rosa, H. T.; Walter, L. C.; Paula, G. M. de; Camera, C.; Lago, I.; Marcolin, E. Evaluation of photoperiodical answer of paddy rice genotypes. *Bragantia*, v.65, p.533-541, 2006. <https://doi.org/10.1590/S0006-87052006000400001>
- Tedesco, M. J.; Gianello, C.; Bissani, C. A.; Bohnen, H.; Volkweiss, S. J. Analysis of soil, plants and other materials. 2.ed. Porto Alegre: Universidade Federal do Rio Grande do Sul, 1995. 147p.
- Walter, L. C.; Rosa, H. T.; Streck, N.A. Simulação do rendimento de grãos de arroz irrigado em cenários de mudanças climáticas. *Pesquisa Agropecuária Brasileira*, v.45, p.1237-1245, 2010. <https://doi.org/10.1590/S0100-204X2010001100002>
- Wang, J.; Lin, X.; Sun, Q.; Jena, K. K. Evaluation of cold tolerance for Japonica Rice varieties from different country. *Advance Journal of Food Science and Technology*, v.5, p.54-56, 2013.
- Wang, Z. F.; Wang, J. F.; Wang, F. H.; Bao, Y. M.; Wu, Y. Y.; Zhang, H. S. Genetic control of germination ability under cold stress in rice. *Rice Science*, v.16, p.173-180. 2009. [https://doi.org/10.1016/S1672-6308\(08\)60076-1](https://doi.org/10.1016/S1672-6308(08)60076-1)
- Weerakoon, W. M. W.; Maruyama, A.; Ohba, K. Impact of humidity on temperature-induced grain sterility in rice (*Oryza sativa* L). *Journal of Agronomy and Crop Science*, v.194, p.135-140, 2008. <https://doi.org/10.1111/j.1439-037X.2008.00293.x>
- Zaffari, G. R.; Scheuermann, K. K.; Marschalek, R.; Medeiros, D. S.; Andrade, A. de. Protocol to produce double haploid plants of rice genotypes from the sub-specie *Indica* by the anther culture. *Plant Cell Culture & Micropropagation*, v.10, p.32-40, 2014.